



Reg. No. L9443 Act. No. 36 of 1947

AnnGro®

1L

AN ADJUVANT/PENETRANT

formulated as an emulsion that enhances the uptake of foliar applied plant nutrients.

'N BEVORDERINGSMIDDEL/ PENETREERDER

emulsie wat die opname van blaar-toegediende plantvoedingstowwe bevorder.



ACTIVE INGREDIENT / AKTIEWE BESTANDEEL

Ethyl esters of fatty acids _____ 25 g/l _____ Etiel-esters van vetsure



BioPher™

REGISTRATION HOLDER / REGISTRASIE HOUER

BioPher (Pty) Ltd | Reg. no: 2008/013129/07

P.O. Box 38910, Langenhovenpark, Bloemfontein, 9330 | www.anngro.com

Batch Number | Date of manufacturing
Lot Nommer | Datum van vervaardiging



GENERAL

- CibuSil™ is water soluble and non-toxic to plants when applied as directed.
- For best results apply CibuSil™ according to recommendations based on plant and/or soil analysis.
- Use only as directed. Use protected gear at all times. Store between 10 and 24°C in well-ventilated area.

DIRECTIONS FOR USE

- Shake container well before use.
- Agitate well in spray tank before administration to crops.
- Apply preferable when conditions for leaf uptake are optimal. Consult manufacturer for more information.
- Apply as a foliar spray to crops or via the irrigation system to the soil.
- Apply tank-mixed product as soon as possible.

APPLICATION RATES

Crop	Dosage rate (kg/ton)	Remark
Row crops	Foliar spray 0.25-0.5 L/ha Via irrigation system 0.5-2 L/ha	Use minimum 200 L water/ha for foliar application. Apply at 3-8 leaf stage and again before flowering if required. Use a 1:400 dilution rate or lower concentration for foliar application.
Vegetable crops	Foliar spray 0.25-1 L/ha Via irrigation system 0.5-2 L/ha	Use minimum 300 L water/ha for foliar application. Apply at 3-8 leaf stage and thereafter once monthly as required. Use a 1:500 dilution rate or lower concentration for foliar application.
Fruit tree crops	Foliar spray 1-3 L/ha Via irrigation system as a root drench 2-4 L/ha (minimum 2 mm/20 000 L water/ha)	Once monthly during season as required. Use a 1:1000 dilution rate or lower concentration for foliar application.

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTION

- CibuSil™ has a high alkalinity and therefore avoid direct contact with skin and eyes. Please do not inhale spray mist.
- Due to the high pH of CibuSil™ always conduct a jar test to confirm compatibility with other products.
- Avoid the use of pH buffering agents. Contact manufacturer for any queries.
- Avoid use with heavy metal containing fertilizers (most of micro-elements), magnesium and calcium.