

# SPRAY-KILL 5

FOR FRUITFLY.

Reg. No.: L4288 Act No. 36 of 1947

A water miscible concentrate for the control of fruit fly on fruit and fruit trees.

IRAC INSECTICIDE GROUP CODE	3
-----------------------------	---

### ACTIVE INGREDIENTS:

Cypermethrin..... 2 g/ℓ  
Piperonyl Butoxide..... 10 g/ℓ  
Tetrachlorvinphos..... 20 g/ℓ

### REGISTRATION HOLDER:

**Effecticide (PTY) LTD**

Co. Reg. No.: 08/06493/07

P.O. Box 38273, Booysens, 2016

Emergency no.: 082 446 8946

### DISTRIBUTOR:

**PROTEK, a division of PE-BEE AGRI (PTY) LTD**

Co. Reg. No.: 2005/036308/07

P.O. Box 72, Heidelberg, 1438

Tel: (011) 812 9800 or 0861 PROTEK (0861 77 68 35)

www.protek.co.za



**CAUTION**



### WARNINGS:

- Allow the following number of days between last application and harvesting or feeding/grazing: 14 days
- Harmful if swallowed.
- Toxic to fish and bees.
- Store away from food and feed.
- Keep out of reach of children and pets.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water; incompatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder, furthermore, does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

### PRECAUTIONS:

- Wash hands with soap and water after use or accidental skin contact.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.
- Dispose of the empty container in a safe manner, according to local regulations and do not re-use for any other purpose.

**DIRECTIONS FOR USE:** Use only as indicated.

### DOSAGE RATES:

CROP:	DOSAGE RATE:	DIRECTIONS FOR APPLICATION:
PEACHES - ALL CULTIVARS	1 ml / 2 ℓ water or 3 ml / 6 ℓ water	Apply as full cover spray 8 weeks before picking and repeat application every 14 days. In summer rainfall regions the first application must be made no later than the third week of December. Repeat sprays every 14 days at high infestations. If infestation is low, repeat spray every 21 days.
EARLY CULTIVARS		Apply as full cover spray 8 weeks before picking and repeat application every 14 days.

Repeat applications as and when necessary.

All applications should be made with suitable equipment that is in good working order.

# SPRAY-KILL 5

VIR VRUGTEVLIEG.

Reg. Nr.: L4288 Wet No. 36 van 1947

'n Water vermengbare konsentraat vir die beheer van vrugtevlieg op vrugte en vrugtebome.

IRAK INSEKDODERGROEP KODE 3

## AKTIEWE BESTANDELE:

Sipermetrien..... 2 g/l  
Piperonielbutoksied..... 10 g/l  
Tetrachloorinfos..... 20 g/l

## REGISTRASIEHOUER:

Effecticide (EDMS) BPK

Mpy. Reg. Nr.: 08/06493/07

Posbus 38273, Booyens, 2016

Nood nr.: 082 446 8946

## VERSPREIDER:

PROTEK, a divisie van PE-BEE AGRI (EDMS) BPK

Mpy. Reg. Nr.: 2005/036308/07

Posbus 72, Heidelberg, 1438

Tel: (011) 812 9800 of 0861 PROTEK (0861 77 68 35)

www.protek.co.za



VERSIGTIG



## WAARSKUWINGS:

- Laat die volgende aantal dae toe tussen laaste bespuiting en oes of voer/beweiding: 14 dae
- Skadelik indieningesluk.
- Giftig vir visse en bye.
- Stoor weg van voedsel en voer.
- Hou buite bereik van kinders en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale gronds-, klimaats- en bergingstoestande; kwaliteit van verdunningswater; onverenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die plaag teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

## VOORSORGMATREËLS:

- Was hande met seep en water na gebruik of toevallige velkontak.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Doen weg met die leë houër op 'n veilige manier, volgens plaaslike regulasies en moet nie hergebruik vir enige ander doel nie.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

## DOSISSE:

GEWAS:	DOSIS:	TOEDIENINGSVOORSKRIFTE:
PERSKES - ALLE VARITEITE	1 ml / 2 l water of 3 ml / 6 l water	Dien toe as 'n dekbespuiting 8 weke voor oes herhaal toediening elke 14 dae. In somerreënval gebiede moet die eerste toediening nie later as die 3de week in Desember gemaak word nie. Herhaal elke 14 dae waar hoë besmettings opgemerk word. Indien besmettings laag is kan toediening elke 21 dae herhaal word.
VROEË VARITEITE		Dien toe as 'n dekbespuiting 8 weke voor oes herhaal toediening elke 14 dae.

Herhaal toediening indien nodig.

Alle toedienings moet gemaak word met geskikte toerusting wat in 'n goeie werkende toestand is.